

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 81300823.2

(51) Int. Cl.³: **C 07 C 121/75**
A 01 N 37/38, C 07 B 19/00

(22) Date of filing: 27.02.81

(30) Priority: 21.04.80 US 142522

(43) Date of publication of application:
28.10.81 Bulletin 81/43

(84) Designated Contracting States:
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: ZOECON CORPORATION
975 California Avenue
Palo Alto California 94303(US)

(72) Inventor: Anderson, Richard James
1021 Metro Circle
Palo Alto California 94303(US)

(72) Inventor: Adams, Karen Gill
7622 Whitehaven Court
West Chester Ohio 45068(US)

(72) Inventor: Henrick, Clive Arthur
3177 Manchester Court
Palo Alto California 94303(US)

(74) Representative: Harrison, David Christopher et al,
MEWBURN ELLIS & CO 70 & 72 Chancery Lane
London WC2A 1AD(GB)

(54) Diastereomers of amino acid esters, pesticidal compositions including them and method of controlling pests using them.

(57) The two specific diastereomers (S)- α -cyano-3-phenoxybenzyl (R)-2-(2-chloro-4-trifluoromethylphenylamino)-3-methylbutanoate and (S)- α -cyano-3-phenoxybenzyl (R)-2-(4-trifluoromethylphenylamino)-3-methylbutanoate are effective for the control of insect or acarid pests. The former compound is preferred.

EP 0 038 617 A1

BEST AVAILABLE COPY


DIASTEREOMERS OF AMINO ACID ESTERS PESTICIDAL
COMPOSITIONS INCLUDING THEM AND METHOD OF
CONTROLLING PESTS USING THEM.

5 This invention relates to certain diastereomers of amino acids esters and to the use thereof for the control of pests.

10 A wide range of esters of substituted-phenylamino acids have been described by Henrick & Garcia, Offenlegungsschrift 28 12 169, as being effective agents for the control of pests such as insects and acarids, acting in the manner of synthetic pyrethroids. These were not stereospecific and the acids and alcohols used were racemic, resulting in
15 racemic products.

20 The diastereomer of formula (A) herein has now been found to possess greatly improved pesticidal activity than (a) any other single specific diastereomer of the same ester (b) any diastereomeric pair of the same ester and (c) any racemic mixture of the same ester.



25 The esters of amino acids of the present invention are represented by the following generic formula (A) :



wherein,

Y is hydrogen or chloro; and
the acid is the R configuration and the
alcohol is the S configuration. Y is chloro gives the
35 preferred ester.

The compounds of formula (A) can be prepared by the reaction of an acid of formula (I), in its R configuration, with the S enantiomer of α -cyano-3-phenoxybenzyl alcohol (II).



The above esterification can be carried out at a low temperature in the presence of 4-dimethylaminopyridine and dicyclohexylcarbodiimide.

The acid (I) is prepared from (R)-valine by conversion first to (R)-2-bromo-3-methylbutanoic acid and reaction of the latter with 4-trifluoromethylaniline to give the 4-substituted phenylamino acid. This can be chlorinated by use of N-chlorosuccinimide to prepare an acid of formula (I) wherein Y is chloro.

The S enantiomer of the alcohol II can be made by reacting racemic α -cyano-3-phenoxybenzyl alcohol with (R)-1-(1-naphthyl)ethyl isocyanate in the presence of 4,4-dimethylaminopyridine and a solvent such as toluene or benzene. The resulting carbamate is separated into its two diastereomers by liquid chromatography. The R,R isomer is further purified by crystallization and the R,S isomer, by repeated chromatography. Alternatively, the diastereomers can be separated from the mixture without initial chromatographic separation by adding a seed crystal of substantially optically pure (R,R)-carbamate and crystallizing out the R,R isomer. The separated

R,S diastereomer, in a solvent such as benzene, is reacted with trichlorosilane and triethylamine, at elevated temperature, to give the resulting S enantiomer of formula II.

5 In the prior art, synthetic pyrethroid esters containing the α -cyano-3-phenoxybenzyl alcohol moiety have been separated into their diastereomers after esterification, rather than by taking the desired acid enantiomer and the desired alcohol
10 enantiomer and then esterifying them to make the diastereomer of the ester compound. Cf. Warnant et al., U.S. Patents 4133826 & 4151195, and Stoutamire, U.S. Patent 4176195.

The two diastereomers of formula A which are
15 the only compounds and diastereomers of the invention are highly active pesticides, particularly against insects and acarids.

In the use of the compounds of formula A for combating insects and acarids for the protection of
20 agricultural crops, for example soybeans, cotton, alfalfa, etc., a compound of formula A usually together with a carrier is applied to the locus in a pesticidally effective amount. The carrier can be liquid or solid and include adjuvants such as
25 wetting agents, dispersing agents and other surface active agents. The compounds of formula A can be used in formulations such as wettable powders, solutions, dusts, granules, emulsifiable concentrates, and the like. Suitable solid carriers include natural
30 and synthetic silicates and clays, carbon or charcoal granules, natural and synthetic resins, waxes, and the like. Suitable liquid carriers include water, aromatic hydrocarbons, alcohols, vegetable and mineral oils, ketones, and the like. The amount of a compound
35 of formula A in the formulation can vary widely, generally within the range of about 0.01 percent to about 90.0 percent, by weight.

The compounds of the present invention are effective on many different insects and on acarids. The compounds are effective control agents for insects such as mosquitoes, flies, aphids, weevils and acarids such as the spider mite and ticks. Among the pests against which the compounds of the present invention are pesticidally effective are insects of the order Lepidoptera, Orthoptera, Heteroptera, Homoptera, Diptera, Coleoptera or Hymenoptera, and acarids of the order Acarina including mites of the family Tetranychidae or Tarsonemidae and ticks such as Ornithodoros.

The compounds of the present invention can be used in combination with pyrethroid synergists and/or other pesticides such as the carbamates, phosphates and insect growth regulators, e.g. propoxur, carbaryl, naled, dichlorvos, methoprene, kinoprene, hydroprene, cyhexatin and resmethrin.

Herein and in the appended claims, the first letter designation refers to the configuration of the acid and the second letter designation refers to the configuration of the alcohol. For example, the diastereomer designated RS refers to ^R acid ^S alcohol.

The following examples are provided to illustrate the practice of the present invention. Temperature is given in degrees Centigrade. RT means room temperature.

"Examples" show preparation and use of the specific diastereomers of the present invention; "Preparations" show preparation of intermediates or comparative compounds.

Preparation 1

To 12.4 g (72.5 mmol) of purified (R)-1-(1-naphthyl)ethylamine in 130 ml of dry toluene is

added, with stirring, gaseous hydrogen chloride for about 30 minutes, during which time an additional 100 ml of toluene is added to facilitate stirring. Phosgene is bubbled into the suspension, at RT, for about 20 minutes and then at reflux for 2 hours. Phosgene addition is stopped and the solution is heated at reflux for another 1.5 hours. The toluene is distilled off at atmospheric pressure and the residue is distilled (short path) at 0.30 mm to yield (R)-1-(1-naphthyl)ethyl isocyanate.

A solution of 11.0 g (56.0 mmol) of (R)-1-(1-naphthyl)ethyl isocyanate, 12.6 g (56.0 mmol) of racemic α -cyano-3-phenoxybenzyl alcohol and 150 mg of 4-dimethylaminopyridine in 75 ml of toluene is heated at 50°, under nitrogen, for about 20 hours. The reaction mixture is cooled and poured into ether and 5% HCl. The organic phase is separated and washed with saturated sodium bicarbonate and with brine and is dried over sodium sulfate. Removal of solvent gives (R,S)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)ethylcarbamate.

25 Preparation 2

The (R,S)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)ethylcarbamate is purified by liquid chromatography on silica columns using ~23% ether/hexane. The first fraction, containing a high ratio of the faster eluting diastereomer, is collected and combined with several ml of ether. Hexane is added until crystals begin to form. This is allowed to crystallize overnight. The resulting crystals are collected and washed with hexane, giving (R)- α -cyano-3-phenoxybenzyl (R)-N-1-

(1-naphthyl)ethylcarbamate, m.p. 121.5-122°, specific rotation $[\alpha]_D^{25} = -15.2^\circ$ (c=10 mg/ml in CHCl_3), diastereomer purity= \sim 99%.

5

Preparation 3

The second fraction obtained from purification of (R,S)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)ethylcarbamate by liquid chromatography, from
10 Example 2 above, which second fraction contains a high ratio of the slower eluting diastereomer, is collected. Since the slower eluting diastereomer does not readily crystallize out of solution, the fraction is further purified by liquid
15 chromatography, using ether/hexane, and collection again of the second fraction. This purification process is continued until a substantially diastereomerically pure example of the compound (S)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)
20 ethylcarbamate is obtained, m.p. 41-41.5°, specific rotation $[\alpha]_D^{25} = -19.6^\circ$ (c=10 mg/ml in CHCl_3), diastereomer purity= \sim 98%.

Preparation 4

25

To 1.99 g (4.7 mmol) of (R)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)ethylcarbamate in 20 ml of benzene is added 725 μ l (525 mg, 5.2 mmol) of triethylamine. The solution, under nitrogen, is
30 stirred while 505 μ l (675 mg, 5.0 mmol) of trichlorosilane is added. The reaction is warmed to 50° for 2.5 hours, and is then poured into saturated ammonium chloride and ether. The organic fraction is washed again with saturated
35 ammonium chloride and then with brine (3X), and is dried over sodium sulfate overnight in the freezer. Solvent is removed by rotoevaporation, and the

residue is washed repeatedly with hexane to remove the isocyanate. The urea contamination is removed by dissolving the product in ether/hexane (~1:1), and the solvent is then removed. Purification by thin layer chromatography (tlc) on silica gel plates developed in 30% ethylacetate/hexane yields the product, which is then dissolved in trichloromethane. The solvent is removed, giving (R)- α -cyano-3-phenoxybenzyl alcohol, specific rotation $[\alpha]_D = +15.2^\circ$ (c=10 mg/ml in acetone).

Preparation 5

To a solution of 2.91 g (6.99 mmol) of (S)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl) ethylcarbamate in 30 ml of dry benzene under a nitrogen atmosphere is added 1.07 ml (0.78 g, 7.7 mmol) of triethylamine followed immediately by 0.75 ml (1.0 g, 7.4 mmol) of trichlorosilane. The reaction is heated at 50° for 3 hours and is then worked up and purified following the procedure of Example 4. The more polar band on the TLC plate is removed and extracted with trichloromethane to yield (S)- α -cyano-3-phenoxybenzyl alcohol, specific rotation $[\alpha]_D^{25} = -14.6^\circ$ (c=11 mg/ml in acetone).

Preparation 6

To a solution of 20.0 g (170 mmol) of (R)-valine in 200 ml of 6 N hydrogen bromide, stirred under nitrogen in an ice bath, is added 18.0 g (260 mmol) of sodium nitrite portionwise over 1.5 hours, keeping the temperature below 8°. On completion of the addition, the reaction is stirred at 5° under nitrogen for about 5 hours and is then

stored in the freezer overnight. Sodium chloride is added to saturate the solution and is then extracted with trichloromethane (3X). The combined organic extracts are washed with sodium bisulfite and with sodium chloride and dried over magnesium sulfate. Evaporation of the solvent under vacuum gives (R)-2-bromo-3-methylbutanoic acid, specific rotation $[\alpha]_D^{25} = +15.5^\circ$ (c=10% in methanol).

A solution of 10.01 g (55.3 mmol) of (R)-2-bromo-3-methylbutanoic acid in methanol is titrated with 1 M potassium hydroxide (~53 ml) in methanol to the phenolphthalein endpoint. The methanol is then removed under high vacuum at 35°, and the solid is kept at 35° under high vacuum for one hour. 4-Trifluoromethylaniline (23.89 g, 148.3 mmol) is added to the salt, and the mixture is heated in a pre-heated oil bath (90°) under nitrogen for 30 minutes. The mixture is then cooled and worked up immediately by partitioning between pentane/5% sodium hydroxide. The layers are separated and the pentane phase is extracted with 5% sodium hydroxide (2X). The combined aqueous phases are extracted with pentane (2X) and acidified with cold conc. hydrochloric acid. The aqueous phase is then extracted with ether (3X), and the combined ether phases are washed with brine and dried over magnesium sulfate. Filtration and evaporation yield (R)-2-(4-trifluoromethylphenylamino)-3-methylbutanoic acid.

To 35 ml of carbon tetrachloride are added 2.81 g (10.0 mmol) of (R)-2-(4-trifluoromethylphenylamino)-3-methylbutanoic acid and 1.33 g (9.96 mmol) of N-chlorosuccinimide, and the mixture is heated to reflux for 1 hour. The mixture is worked up by partition between ether/water. The layers are separated and the aqueous phase is extracted with

ether (2X). The combined organic extracts are washed with water and with brine, dried over magnesium sulfate, filtered and evaporated, giving (R)-2-(2-chloro-4-trifluoromethylphenylamino)-3-methylbutanoic acid.

Preparation 7

(S)-2-bromo-3-methylbutanoic acid is prepared following the procedure of Preparation 6 from (S)-valine and sodium nitrite and 6 N hydrogen bromide. Specific rotation of the bromo acid is $[\alpha]_D^{25} = -17.2^\circ$ (c= 10% in methanol).

1.5 Grams (8.3 mmol) of the (S)-2-bromo-3-methylbutanoic acid is titrated with 8 ml of 1 M potassium hydroxide in methanol to a phenolphthalein endpoint. The solution is evaporated under high vacuum at 40° for 1 hour, and 0.51 g (2.33 mmol) of the resulting salt is heated together with 1.52 g (9.43 mmol) of 4-trifluoromethylaniline at 90-95° under nitrogen for 1 hour. Heating is discontinued and the reaction worked up immediately by partition between ether/5% sodium hydroxide. The layers are separated and the organic phase is extracted with 5% sodium hydroxide (2X). The aqueous layers are combined and washed with ether (2X), then acidified with ice and conc. HCl and extracted with ether (3X). The latter ether extracts are washed with brine, dried over magnesium sulfate, filtered and evaporated, giving (S)-2-(4-trifluoromethylphenylamino)-3-methylbutanoic acid.

Following the method of Preparation 6, (S)-2-(4-trifluoromethylphenylamino)-3-methylbutanoic acid and N-chlorosuccinimide in carbon tetrachloride are reacted, yielding (S)-2-(2-chloro-4-

trifluoromethylphenylamino)-3-methylbutanoic acid.

Example 1

5 Into 4.5 ml of methylene chloride containing
22.5 mg of 4-dimethylaminopyridine are added 439 mg
(1.48 mmol) of (R)-2-(2-chloro-4-
trifluoromethylphenylamino)-3-methylbutanoic acid and
329 mg (1.46 mmol) of (S)- α -cyano-3-phenoxybenzyl
10 alcohol. The solution is cooled in an ice bath
and 380 mg (1.82 mmol) of dicyclohexylcarbodiimide is
added. The mixture is stirred at 0° for 1 hour
and is then worked up by partition between ether/
hexane. The aqueous phase is extracted with ether,
15 and the ether extract is washed with water (2X)
and brine, dried over magnesium sulfate, filtered
and evaporated. The product is purified by radial
thin layer chromatography (2mm silica gel rotor
eluting with 12% ether/hexane). Higher
20 diastereomeric purity is obtained by further
chromatron tlc, yielding the final product (S)- α -
cyano-3-phenoxybenzyl (R)-2-(2-chloro-4-
trifluoromethylphenylamino)-3-methylbutanoate,
specific rotation = +46.3° (in CHCl₃), diastereomer
25 purity = ~98%..

Preparation 8

30 (R)-2-(2-chloro-4-trifluoromethylphenylamino)-3-
methylbutanoic acid (0.20 g, 0.68 mmol), racemic
 α -cyano-3-phenoxybenzyl alcohol (0.17 g, 0.75 mmol)
and dimethylaminopyridine (10 mg, 0.08 mmol) are
placed in 2 ml methylene chloride/2ml dimethyl-
formamide. The mixture is cooled in an ice bath
35 under nitrogen and 0.16 g (0.77 mmol) of

- 11 -

dicyclohexylcarbodiimide is added. The mixture is kept cold, with stirring, for 2 hours.

5 The mixture is filtered and the solid is washed several times with ether. The ether is then extracted with water and the layers are separated. The aqueous phase is extracted with ether (2X), and the combined ether extracts are washed with dilute HCl and with brine, dried over magnesium sulfate, filtered and evaporated.

10 Purification by preparative thin layer chromatography eluting with 10% ether/hexane gives a product composed primarily of the diastereomer pair consisting of (S)- α -cyano-3-phenoxybenzyl (R)-2-(2-chloro-4-trifluoromethylphenylamino)-3-

15 methylbutanoate (42%) and (R)- α -cyano-3-phenoxybenzyl (R)-2-(2-chloro-4-trifluoromethylphenylamino)-3-methylbutanoate (52%).

Preparation 9

20 Approximately 0.5 g of the compound (R,S)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)ethylcarbamate from Preparation 1 is dissolved in ~1 ml of ether and a few drops of hexane are

25 added. To this is added a seed crystal of ~99% optically pure (R)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)ethylcarbamate from Example 2, and the mixture is allowed to crystallize over night. The solid is separated out and washed with hexane

30 several times. The solid shows a diastereomer purity of 72% (R)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)ethylcarbamate. Several additional crystallizations of this solid gives compound of 98% diastereomeric purity.

35 The separated mother liquor is purified by

BAD ORIGINAL

liquid chromatography as in Preparation 3 to give
(S)- α -cyano-3-phenoxybenzyl (R)-N-1-(1-naphthyl)
carbamate.

5

Example 2

Following the procedure of Example 1,
(R)-2-(4-trifluoromethylphenylamino)-3-methylbutanoic
acid and (S)- α -cyano-3-phenoxybenzyl alcohol are
10 combined, in the presence of 4-dimethylaminopyridine
and dicyclohexylcarbodiimide, to yield (S)- α -cyano-
3-phenoxybenzyl (R)-2-(4-trifluoromethylphenylamino)-
3-methylbutanoate, specific rotation = +62.1°
(c=9 mg/ml in CHCl₃).

15

Example 3

Comparative activity of the four diastereomers
of α -cyano-3-phenoxybenzyl 2-(2-chloro-4-
20 trifluoromethylphenylamino)-3-methylbutanoate and
of the prior art diastereomeric mixture of the
compound was determined by testing for toxicity
on insect pests.

A. Two groups of 10 each of 0-24 hr III
25 instar Heliothis virescens larvae were treated with
1 μ l of the test compound in acetone at different
dosage rates by application to the dorsum of the
thorax. Two groups of 10 each were treated
identically with 1 μ l acetone as controls. Larvae
30 are held individually in 30 ml plastic cups provided
with artificial medium for 72 hours at 25° and
16 hr photoperiod. After 72 hr the number of dead
is calculated as a percentage of the total number
originally treated and then corrected for any
35 mortality in the control group using Abbott's formula.

The toxicity is expressed as LD₅₀, which is the dosage, in µg per insect, required to kill 50% of the test insects. The results are presented in Table I.

- 5 B. Fifteen 72-hr-old adult female Musca domestica L. are anaesthetized with ether vapor. These are then treated with 1 µl of the test compound diluted to different dosage rates in acetone applied to the dorsal surface of the
- 10 prothorax. They are held in an assay container with milk-saturated cotton at 25°, 16 hr photoperiod for 24 hours. The effect is stated as the number dead calculated as a percentage of the total, corrected for any control mortality using
- 15 Abbott's formula. The toxicities of the compounds, expressed as LD₅₀, are presented in Table 1 below.

Table I

20

ACTIVITY OF α-CYANO-3-PHENOXYBENZYL
2-(2-CHLORO-4-TRIFLUOROMETHYLPHENYLAMINO-3-METHYLBUTANOATE
COMPOUNDS, AS LD₅₀ (µg/INSECT)

25

Compound	Diastereomer	<u>H. virescens</u>	<u>M. domestica</u>
A	<u>RR</u>	0.320	2.180
B	<u>SR</u>	91.000	29.900
30 C	<u>SS</u>	>100	39.900
D*	<u>RS</u>	0.0181	0.0378
E	<u>RS,SR,RR,SS</u>	0.0803	0.156

* Compound of the present invention

35



The results of the above tests show that compound D, (S)- α -cyano-3-phenoxybenzyl (R)-2-(2-chloro-4-trifluoromethylphenylamino)-3-methylbutanoate, is more than four times as active as the diastereomeric mixture (R,S)- α -cyano-3-phenoxybenzyl (R,S)-2-(2-chloro-4-trifluoromethylphenylamino)-3-methylbutanoate (compound E). The three other diastereomers are all much less active than the diastereomeric mixture (compound E).

Example 4

Following the methods described in Example 3, comparative activity of the four diastereomers and the diastereomeric mixture of α -cyano-3-phenoxybenzyl 2-(4-trifluoromethylphenylamino)-3-methylbutanoate against III instar Heliothis virescens larvae and against adult female Musca domestica was determined. The results are presented in Table II.

Table II


ACTIVITY OF α -CYANO-3-PHENOXYBENZYL 2-(4-TRIFLUOROMETHYLPHENYLAMINO)-3-METHYLBUTANOATE. COMPOUNDS, AS LD ₅₀ (μ g/INSECT)			
Compound	Diastereomer	<u>H. virescens</u>	<u>M. domestica</u>
F	<u>RR</u>	0.53	0.82
G	<u>SR</u>	>100	2.28
H	<u>SS</u>	2.53	0.17
J*	<u>RS</u>	0.018	0.033
K	<u>RS,SR,RR,SS</u>	0.055	0.084
*Compound of the present invention			

Compound J, (S)- α -cyano-3-phenoxybenzyl
(R)-2-(4-trifluoromethylphenylamino)-3-
methylbutanoate, is three times as active as the
diastereomeric mixture (R,S)- α -cyano-3-

- 5 phenoxybenzyl (R,S)-2-(4-trifluoromethylphenylamino)-3-
methylbutanoate (compound K) on Heliothis and more
than 2.5 times as active on Musca. The other three
diastereomers have much lower activity.

CLAIMS:

1. A compound of the following Formula A:



- 10 Y is hydrogen or chloro; and
the acid is the R configuration and the
alcohol is the S configuration.

- 15 2. The diastereomer (S)- α -cyano-3-
phenoxybenzyl (R)-2-(2-chloro-4-
trifluoromethylphenylamino)-3-methylbutanoate.

- 20 3. The diastereomer (S)- α -cyano-3-
phenoxybenzyl (R)-2-(4-trifluoromethylphenylamino)-3-
methylbutanoate.

- 25 4. A method for controlling insects or acarids
which comprises applying to said insect or acarid or
their habitat a compound as claimed in claim 1,
claim 2 or claim 3.

5. A pesticidal composition comprising as active
ingredient a compound as claimed in claim 1, claim 2
or claim 3.



European Patent
Office

EUROPEAN SEARCH REPORT

0038617

Application number

EP 81 30 0823

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
D	<u>DE - A - 2 812 169</u> (ZOECON CORP.) * Page 4: 5 ^o , page 20 * & GB - A - 1 588 111 --	1,4	C 07 C 121/75 A 01 N 37/38 C 07 B 19/00
	<u>EP - A - 0 002 289</u> (SHELL) * Pages 1,2 * ----	1,4	TECHNICAL FIELDS SEARCHED (Int. Cl.) C 07 C 121/75 A 01 N 37/38 C 07 B 19/00
			CATEGORY OF CITED DOCUMENTS
			X: particularly relevant A: technological background O: non-written disclosure P: intermediate document T: theory or principle underlying the invention E: conflicting application D: document cited in the application L: citation for other reasons
			&. member of the same patent family, corresponding document
<input checked="" type="checkbox"/> The present search report has been drawn up for all claims			
Place of search	Date of completion of the search	Examiner	
The Hague	08-07-1981	VERHULST	

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☐ BLACK BORDERS

☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES

☐ FADED TEXT OR DRAWING

☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING

☐ SKEWED/SLANTED IMAGES

☒ COLOR OR BLACK AND WHITE PHOTOGRAPHS

☒ GRAY SCALE DOCUMENTS

☒ LINES OR MARKS ON ORIGINAL DOCUMENT

☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY

☐ OTHER: _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.